

Hernia: Inguinal – Surgical anatomy, presentation, treatment, complications

**Dr S. Gulve
Associate Professor
Dept Of Surgery**

Introduction

Abnormal protrusion of viscus or a part of it
through a weak point in the abdominal wall


Anatomy of inguinal region

- **Superficial inguinal ring-**
 - triangular aperture in the aponeurosis of the ext oblique muscle .
 - Lies 1.25 cm above the pubic tubercle .
 - Normally it doesn't admit the tip of the little finger.
- **Deep inguinal ring –**
 - U shaped condensation of the fascia trasversalis
 - Lies 1.25cm above the mid inguinal point.

Inguinal canal

- Oblique passage in the lower part of the anterior abdominal wall.
- Extends from deep inguinal ring to superficial inguinal ring.
- Directed downwards forwards and medially
- About 4cm long

Small bowel



Boundaries

- Anterior – Ext. oblique aponeurosis & conjoined muscle laterally.
- Posterior – Fascia transversalis & the conjoined tendon.
- Superiorly – conjoined muscle.
- Inferiorly – inguinal ligament.

Contents

- Spermatic cord
- Ilioinguinal nerve
- Genital branch of genitofemoral nerve
- Females – Round ligament is present instead of spermatic cord.

Spermatic cord constitutes- vas deferens, testicular & cremastic arteries , pampiniform plexus of veins, lymphatics

Predisposing

- All hernias occur at the site of WEAKNESS OF THE ABDOMINAL WALL which are acted on by repeated INCREASE in abdominal pressure

repeated INCREASE in abdominal pressure is usually due to

- ●Chronic cough
- ●Straining
- ●Bladder neck or urethral obstruction
- ●Pregnancy
- ●Vomiting
- ●Sever muscular effort
- ●Ascetic fluid

TYPES

- Inguinal
- Femoral
- Epigastric
- Para umbilical
- Umbilical
- Obturator
 - Superior lumbar
- Inferior lumbar
 - Gluteal
- Sciatic
- Incisional

- • Indirect Inguinal Hernia -Hernia through the inguinal canal
- • Direct Inguinal Hernia- The sac passes through a weakness or defect of the transversalis fascia in the posterior wall of the inguinal canal
- • Femoral Hernia -Hernia medial to femoral vessels under inguinal ligament
- • Umbilical Hernia- Hernia through the umbilical ring
- • Paraumbilical Hernia A protrusion through the linea alba just above or sometimes just below the umbilicus
- • Epigastric Hernia- Protrusion of extraperitoneal fat through the linea alba anywhere between the xiphoid process and the umbilicus
- • Incisional Hernia- Hernia through an incisional site
- • Lumbar Hernia occur through the inferior lumbar triangle of Petit



Male inguinal hernia



Female inguinal hernia

Clinical types

- **Reducible** –contents can be returned into the abdominal cavity.
- **Irreducible** – contents cannot be returned into the abdominal cavity.
- **Obstructed** – irreducibility + intestinal obstruction, but the blood supply is not impaired.
- **Strangulated**- irreducibility + intestinal obstruction+ arrest of the blood supply.
- **Inflamed**- rare condition. Occurs when contents eg. Appendix,meckel's diverticulum is inflamed

Epidemiology

- **Approximately 7% of all surgical outpatient.**
- Accounts for 96% groin hernias (other 4% are femoral)
- Bilateral in 20% of cases
- Lifetime risk of inguinal hernia: 10%
- M:F 9:1

- Affects 1-3% of young children
- In men the incidence rises from 11 per 10,000 person years aged 16-24 years to 200 per 10,000 person years aged 75 years or above.
- Extremely common; represents the most frequent problem requiring surgical intervention in the paediatric age group
- Much more common in boys (90% of cases) than girls
- Definite familial tendency,
- more frequent on the right side as a result of later descent of the right testis and delayed obliteration of the right processus vaginalis.

Presentation

- Pain
 - Localized pain
 - Referred pain
 - Generalized pain
- Nausea and vomiting
- Constipation
- Urinary symptoms

Presentation

- At first appearance, it is easily reducible.
- With time it can no longer be reduced, it is irreducible or incarcerated.
- Strangulation: when visceral contents of the hernia become twisted or entrapped by the narrow opening.


Strangulation usually leads to bowel obstruction with sudden, severe pain in the hernia, vomiting and irreducibility.

Nyhus Classification System

Type I	Indirect hernia; internal abdominal ring normal; typically in infants, children, small adults
Type II	Indirect hernia; internal ring enlarged without impingement on the floor of the inguinal canal; does not extend to the scrotum
Type IIIA	Direct hernia; size is not taken into account
Type IIIB	Indirect hernia that has enlarged enough to encroach upon the posterior inguinal wall; indirect sliding or scrotal hernias are usually placed in this category because they are commonly associated with extension to the direct space; also includes pantaloon hernias
Type IIIC	Femoral hernia
Type IV	Recurrent hernia; modifiers A-D are sometimes added, which correspond to indirect, direct, femoral, and mixed, respectively

Diagnosis- Inspection

- Inguinal hernias are best examined with the patient standing.
- Coughing may increase the size of the hernia.
- Site and shape of the hernia:
 - those appearing above and medial to the pubic tubercle are inguinal hernias
 - those appearing below and lateral to the pubic tubercle are femoral hernias
- whether the lump extends down into the scrotum
- any other scrotal swellings
- any swellings on the 'normal' side
- scar from previous surgery or trauma



Digital examination of the inguinal canal

Palpation

- Confirm inspectory findings
- Examine the scrotum- Getting above the swelling is not possible
- Consistency, temperature, tenderness and fluctuance.
- One should attempt to reduce the hernia: Ask the patient to reduce. Otherwise flex and medially rotate the hip and reduce
- If the hernia cannot be reduced the probable identity of the hernia is: femoral > indirect inguinal > direct inguinal
- Expansile cough impulse

- **Deep ring occlusion test-** reduce the swelling
- Locate the deep ring 1/2 “ above the midpoint of the inguinal ligament and occlude it asking the patient to cough.
- Impulse seen- direct, not seen- indirect
- Leg raising test- Malgaigne’s bulgings seen
- **Zieman’s method**
- Swelling gurgles- enterocoele, firm/granular- omentocoele.
- Always palpate the other inguino-femoral region as herniae are often bilateral

Percussion

The characteristics of hernias depend on their contents:

- bowel is hyper-resonant and has bowel sounds unless it is strangulated
- omentum and fat is dull and does not have bowel sounds

Investigations

Ultrasound

- High [Test Sensitivity](#) (>90%)
- High [Test Specificity](#)
 - Distinguish [Incarcerated Hernia](#) from firm mass

Herniography

- Suspected hernia, but clinical dx unclear
- Procedure done under flouroscopy following injection of contrast medium
- Frontal and oblique radiographs are taken with and without increased intra-abdominal pressure

Systemic examination

- Examine respiratory system
- Per rectal examination
- Abdominal
- Ext genitalia

Complications

Bowel incarceration (acute, chronic): The trapping of abdominal contents within the *Hernia* itself

Strangulation: **pressure** on the hernial contents may compromise blood supply (especially veins, with their low pressure, are sensitive, and venous congestion often results) and cause ischemia, and later necrosis and gangrene, which may become fatal.

Small Bowel Obstruction

Management

Non operative Treatment

- Watchful waiting: for asymptomatic or minimally symptomatic

Truss is a mechanical appliance ,belt with a pad applied to groin after spontaneous or manual reduction of hernia

The purpose is twofold: to maintain reduction and to prevent enlargement.

Surgery

Mesh repairs

Open repair (Lichtenstein, Shouldice, Bassini)

Most commonly performed: Lichtenstein repair


It's "tension-free" repair

Tension-free repairs

- Desarda
- Guarnieri

Bassini technique, first suture:

- Aponeurosis musculi obliq. ext.
- Musculus obliquus internus
- Musculus transversalis
- Fascia transversalis
- Peritoneum
- Ligamentum inguinale.



Laparoscopic repair

- transabdominal preperitoneal (TAPP)
- totally extra-peritoneal (TEP) repair

- **THANK YOU**